

Aberystwyth University

New ExoMol line lists for PO, PS, SiH, NS and SH.

Gorman, Maire; Yurchenko, Sergei N.; Tennyson, Jonathan; Bond, Wesley; Hill, Christian; Jagoda, Pawel; Lodi, Lorenzo; McKemmish, Laura K.; Nunn, William; Prajapat, Laxmi; Sinden, Frances

Publication date:
2018

Citation for published version (APA):

Gorman, M., Yurchenko, S. N., Tennyson, J., Bond, W., Hill, C., Jagoda, P., Lodi, L., McKemmish, L. K., Nunn, W., Prajapat, L., & Sinden, F. (2018). *New ExoMol line lists for PO, PS, SiH, NS and SH.* Poster session presented at EWASS/NAM 2018, Liverpool, United Kingdom of Great Britain and Northern Ireland.

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Maire N. Gorman^{1, 2}, Sergey N. Yurchenko¹, Jonathan Tennyson¹, Wesley Bond¹, Christian Hill¹, Pawel Jagoda¹, Lorenzo Lodi¹, William Nunn¹, Laura McKemmish^{1, 3}, Laxmi Prajapat¹, Rohan Shah¹, Frances Sinden¹

1. Department of Physics and Astronomy, University College London, London WC1E 6BT, UK
2. Department of Physics, Aberystwyth University, Penglais, Aberystwyth, Ceredigion, UK, SY23 3BZ
3. School of Chemistry, University of New South Wales, Sydney, Australia, NSW 2052

Introduction

In this poster we present recently published line lists for five diatomic molecules which are of astronomical interest as detailed below:

SiH

- Detected in sunspots (Pearse 1933 and Babcock 1945) and Solar disk spectrum (Schadee 1964 and Moore-Sitterly 1966, Lambert & Mallia 1970, Grevesse & Sauval 1971).
- Detected in late type stars (Davis 1940) and M- and S-type Mira variable stars (Merrill 1955).
- Of interest in M-dwarf atmospheres (Allard & Hauschildt 1995).
- Prediction that should be present in exoplanetary & brown dwarf atmospheres (Visscher et al. 2010).
- Candidate for ISM clouds (Herbst et al. 1989).

SH

- Challenging to detect due to location of key rotational transition.
- Detection in AGB stars (Yamamura et al. 2000) & Sun's atmosphere (Berdyugina & Livingston 2002).
- Tentatively detected in comets (Krishna Swamy & Wallis 1987, 1988).
- Following extensive searches, Neufeld et al. (2012) detected SH in the ISM using SOFIA.
- Predicted to occur in brown dwarfs (Visscher et al. 2006) and hot Jupiter exoplanets (Visscher et al. 2006; Zahnle et al. 2009).

NS

- One of first ten diatomic molecules to be detected in space (Somerville 1977; Lovas et al. 1979).
- Detected in giant molecular clouds (McGonagle et al. 1992; Leurini et al. 2006; Belloche et al. 2013), cold dark clouds (McGonagle et al. 1994), comets (Irvine et al. 1999; Biver 2005), extragalactically (Martín et al. 2003) and the NGC 253 starburst region (Meier et al. 2015).

PO

Detected in the following:

- Red Supergiant Star VY Canis Majoris (Tenenbaum et al. 2007).
- Oxygen-rich AGB star IK Tauri (De Beck et al. 2013).
- Star-forming regions (Lefloch et al. 2016; Rivilla et al. 2016).

PS

- No detection in space yet.
- A systematic attempt at its astronomical detection was performed by Ohishi et al. (1988).
- Local thermodynamic equilibrium calculations by Tsuji (1973) indicate that PS should be the major P-bearing molecule in oxygen-rich circumstellar envelopes for temperatures below 2000 K.

Fig 1. Potential energy curves of SiH used in the line list production. The $X^2\Pi$ and $A^2\Delta$ PECs have been refined, the $a^4\Sigma^-$ is *ab initio* and the $B^2\Sigma^-$ PEC is an artificial object used to improve the description of the Λ -doubling in the X-state spectra.




Fig 2. Comparison of simulated spectra using the new ExoMol line list for ^{32}SH with the CDMS database for the $v = 0$ band.




Fig 3. PO absorption spectrum at $T = 300$ (bottom), 1000, 2000 and 3000 (top) K, presented with cross-sections on a logarithmic scale. A Gaussian profile with $\text{HWHM} = 10 \text{ cm}^{-1}$ was used.




Fig 4. $T = 2000 \text{ K}$ absorption spectra of 28 SiH: X – X, a–X, and A – X bands, where the a – X band is dipole forbidden.




Table 1. Overview of line lists calculated. Here the number of states and transitions is quoted to the nearest 100.

Isotopologues	Jmax	Energies	Transitions
^{28}SiH , ^{29}SiH , ^{30}SiH	82.5	11 800	1 726 000
^{28}SiD	113.5	21 200	3 521 000
^{32}SH , ^{33}SH , ^{34}SH , ^{36}SH	60.5	2 300	81 300
^{32}SD	84.5	4 500	219 500
$^{14}\text{N}^{32}\text{S}$, $^{14}\text{N}^{33}\text{S}$, $^{14}\text{N}^{34}\text{S}$, $^{14}\text{N}^{36}\text{S}$	235.5	31 800	3 331 000
$^{15}\text{N}^{32}\text{S}$	240.5	33 100	3 479 000
$^{31}\text{P}^{16}\text{O}$	234.5	43 100	2 096 000
$^{31}\text{P}^{32}\text{S}$	320.5	226 00	30 395 00

Fig 5. Comparison of spectra (298 K) for the fundamental $v = 1 - 0$ band of NS for $^{15}\text{N}^{32}\text{S}$ against $^{14}\text{N}^{32}\text{S}$.

This work was supported by the UK Science and Technology Research Council (STFC) No. ST/M001334/1 and the COST action MOLIM No. CM1405. This work made extensive use of UCL's Legion high performance computing facility.

